

**Calculus II**  
**Practice Problems 6: Answers**

Determine whether or not the integral converges. If it does, try to find its value (you may not be able to do this in some cases).

1.  $\int_2^\infty \frac{dx}{x(\ln x)^2} =$

**Answer.** This integral converges. We make the substitution  $u = \ln x$ ,  $du = dx/x$ . We get

$$\int_2^a \frac{dx}{x(\ln x)^2} = \int_{\ln 2}^{\ln a} \frac{du}{u^2} = -u^{-1} \Big|_{\ln 2}^{\ln a} = \frac{1}{\ln 2} - \frac{1}{\ln a} \rightarrow \frac{1}{\ln 2}$$

as  $a \rightarrow \infty$ .

---

2.  $\int_1^{10} \frac{dx}{x\sqrt{\ln x}} =$

**Answer.** Since  $\ln 1 = 0$ , the problem is at the lower limit of integration. Nevertheless, this integral converges. We make the same substitution  $u = \ln x$  and get (taking  $a > 0$  and small)

$$\int_a^{10} \frac{dx}{x\sqrt{\ln x}} = \int_{\ln a}^{\ln 10} \frac{du}{u^{1/2}} = 2u^{1/2} \Big|_{\ln 2}^{\ln 10} = 2\sqrt{\ln 10} - 2\sqrt{\ln a} \rightarrow 2\sqrt{\ln 10}$$

as  $a \rightarrow 1$ .

---

3.  $\int_{1/5}^\infty \frac{\ln(5x)}{x^2} dx =$

**Answer.** Since  $\ln x < \sqrt{x}$  for  $x$  large enough, the integrand is less than  $5x^{-3/2}$ , so by comparison (proposition 8.6), our integral converges. We can now proceed to evaluate it. We make the substitution  $u = \ln(5x)$ ,  $du = dx/x$ , and  $x = e^{-u}/5$ :

$$\int_{1/5}^a \frac{\ln(5x)}{x^2} dx = 5 \int_0^{\ln(5a)} e^{-u} u du = 5(e^{-u} u - e^{-u}) \Big|_0^{\ln(5a)} = 5\left(\frac{\ln(5a)}{5a} - \frac{1}{5a} + 1\right)$$

which converges to 5 as  $a \rightarrow \infty$ .

---

4.  $\int_{-\infty}^\infty \frac{dx}{(1+x^2)^{3/2}} =$

**Answer.** This integral converges, by the comparison test. For,  $(1+x^2)^{3/2} \geq 1+x^2$ , so

$$\frac{1}{(1+x^2)^{3/2}} \leq \frac{1}{1+x^2},$$

and the integral of the latter is finite ( $= \pi$ ). Now, we can find the value by a trigonometric substitution. Let  $x = \tan u$ ,  $dx = \sec^2 u du$ ,  $\sqrt{1+x^2} = \sec u$ . (Draw the triangle corresponding to these substitutions!). This gives us

$$\int \frac{dx}{(1+x^2)^{3/2}} = \int \frac{\sec^2 u du}{\sec^3 u} = \int \cos u du = \sin u + C = \frac{x}{\sqrt{1+x^2}} + C.$$

Thus

$$\int_{-a}^b \frac{dx}{(1+x^2)^{3/2}} = \frac{x}{\sqrt{1+x^2}} \Big|_{-a}^b = \frac{b}{\sqrt{1+b^2}} - \frac{-a}{\sqrt{1+a^2}} = \frac{b}{\sqrt{1+b^2}} + \frac{a}{\sqrt{1+a^2}}.$$

Now the limit of this, as  $a$  and  $b$  go to infinity is (as we saw in problem 10 of practice set 5)  $1+1=2$ .

---



---

$$5. \int_0^{\pi/2} \frac{dx}{1-\cos x} =$$

**Answer.** The improprieness here is at  $x=0$  ( $\cos(0)=1$ ). There is no easy comparison to make, so we calculate the integral. To do so we use some trigonometric identities:

$$\frac{1}{1-\cos x} = \frac{1}{1-\cos x} \frac{1+\cos x}{1+\cos x} = \frac{1+\cos x}{1-\cos^2 x} = \frac{1+\cos x}{\sin^2 x}.$$

Thus

$$\int \frac{dx}{1-\cos x} = \int \csc^2 x dx + \int \cot x \csc x dx = -\cot x - \csc x.$$

Now we calculate

$$\int_a^{\pi/2} \frac{dx}{1-\cos x} = -\cot(\pi/2) - \csc(\pi/2) + (\cot a + \csc a) = -1 + (\cot a + \csc a).$$

If we let  $a \rightarrow 0^+$ , the term in parentheses becomes infinite. Thus the integral diverges.

---



---

$$6. \int_0^1 \frac{dx}{(1-x)^{3/2}} =$$

**Answer.** Make the change of variable  $u = (1-x)$ ,  $du = -dx$ . Then the integral becomes  $\int_0^1 u^{-3/2} du$ . But we saw (see display (7) of the Notes), this integral diverges.

---

$$7. \int_0^{1/2} \frac{dx}{\sqrt{x}(1-x)}$$

**Answer.** In this range  $1-x \geq 1/2$ , so

$$\frac{1}{\sqrt{x}(1-x)} \leq \frac{2}{\sqrt{x}},$$

so by comparison the integral converges.

---



---

8. Find the area under the curve  $y = (x^2 - x)^{-1}$ , above the  $x$ -axis and to the right of the line  $x = 2$ .

**Answer.** First, we find the indefinite integral. We find the partial fraction representation

$$\frac{1}{x^2 - x} = \frac{1}{x-1} - \frac{1}{x}.$$

$$\int \frac{dx}{x^2 - x} = \int \frac{dx}{x-1} - \int \frac{dx}{x} = \ln\left(\frac{x-1}{x}\right).$$

Now,

$$\int_2^a \frac{dx}{x^2 - x} = \ln\left(\frac{x-1}{x}\right) \Big|_2^a = \ln\left(\frac{a-1}{a}\right) - \ln\left(\frac{2-1}{2}\right) = \ln\left(\frac{a-1}{a}\right) + \ln 2.$$

As  $a \rightarrow \infty$ , the first term goes to zero, so the answer is  $\ln 2$ .