

Math 6510 - Homework 4

Due in class on 10/3/13

1. Let M be an n -dimensional manifold embedded in \mathbb{R}^m . If $I = \{i_1, \dots, i_n\} \subset \{1, \dots, m\}$ define $\pi_I : \mathbb{R}^m \rightarrow \mathbb{R}^n$ by $\pi_I(x_1, \dots, x_m) = (x_{i_1}, \dots, x_{i_n})$. Show that for all $x \in M$ there exists an I and an neighborhood of U of x in M such that (U, π_I) is a chart.
2. Let M be an n -dimensional manifold embedded in \mathbb{R}^m . Show that for any $k \geq m - n$ there exists a k -dimensional hyperplane P such that $M \cap P$ is a non-empty smooth manifold of dimension of $n + k - m$. If $k < m - n$ show that there exists k -dimensional hyperplane that is disjoint from M .
3. Let $f : M \rightarrow M$ be a smooth map and $x \in M$ a fixed point ($f(x) = x$). Show that if 1 is not an eigenvalue of $f_*(x)$ that x has a neighborhood U such that if $y \in U$ and $f(y) = y$ then $y = x$. (Hint: Look at the graph of f in $M \times M$ and show that it intersects the diagonal transversally at (x, x) .)